DIVISION FOR PLANETARY SCIENCE

ICES ABSTRACT FO	RM —			—+
	Ru	n.No	Sess.No	
		FOR 5	EDITORIAL USE ONLY	
ORA'L PAPER XX		POSTER	PAPER	
PAPER PRESENTED BY	' N. RAPPA	PORT		
	(P'ease	Print, Must be	First Author)	
SPECIAL INSTRUCTION	10.			
SPECIALINSTRUCTION	(8)			
JPL 301-150				
First Author's Address – 4800 Oak Grove Dri	Print	Signatur	e of First Author	
Pasadena, CA 91109				
LEGGLESIA, OH JEEO			e of Introducing Membe is a Non-member	۲,
E-Vail NUB%NOYAD@HAY	<u> </u>	hone(818) 354-8211	
NUB%NOMAD@HAM	LET.CALTECH	.EDU		
Membership Status (Pre	senter):			
DPS-AAS Member X	Non-m	ember	Student	
ls your abstract newswo publicity staff with addition	rthy, and if so v ona! materia! or	vould you b	e willing to assist ou for reporters.	ır
Yes	No X	Mayb	e	
DPS Category No. 7	(From list on s	separate pa	ae)	
Abstracts must conform	•		0 /	r
MOSCACIS INUSCICIONATORMO	TO THE ALS STATE	שב ה שב היוי	コログ ひわ ちりひ わりんじ ヘチナ	nin ti

Abstract originals must be typed or glued directly onto this form. The charge for publication of this abstract in the Bulletin of the American Astronomical

DPS Abstracts

Society will be included in the registration fee for this meeting.

Deadline for receipt of abstract: July 19, 1993.

SUBMIT ORIGINAL AND FIVE COPIES TO:

Viscous Excitation of Coupled Librations Modes in Planetary Rings

N. Rappaport (ex Borderies) (JPL), P.-Y. Longaretti (Grenoble Obs.)

Viscous instabilitites can develop in a dense ring (Borderies, Goldreich and Tremaine, 1985, Icarus 63, 406-420). An analysis of viscous instabilities based on a two streamlines model shows that, within this simplified model, two regimes of instability exist. In the first regime, librations of the streamlines are excited and persist whereas the mean eccentricity of the ring is damped. In the second regime, the librations are damped whereas the mean eccentricity is driven to an equilibrium value depending on the details of the stress tensor model under consideration. The two regimes are differentiated by the value of the initial eccentricity.

Viscous excitation of coupled libration modes may be relevant to a number of unsolved problems in planetary rings. Examples are the rigid precession of non-circular and non-equatorial narrow rings, the unexplained large residuals of the a ring of Uranus, and the origin of the small scale structures of the Bring of Saturn.

We have written a program that allows us to study the dynamical evolution of streamlines in the presence of a superposition of libration modes. We will present preliminary results concerning the various instability regimes and the temporal and spatial behavior of the libration modes.

Abstract Submitted for the Division for Planetary Sciences Meeting, Boulder, CO, USA

Date Submitted

Form Version 2/93

FOR EDITORIAL USE ONLY

Lunar and Planetary Institute **Publications Services Department**

3600 Bay Area Boulevard Houston, TX 77058-1113